SPECIALTY INFUSION CENTER
An Access Infusion Care Company

ORENCIA (ABATACEPT) ORDERS

PATIENT INFORMATION

Name: _____ DOB: _____ Height: _____ Weight: _____

Address: _____ Phone: _____ Allergies: _____

City, State, Zip: _____ Email: _____

REFERRAL STATUS

☐ New Referral ☐ Dose or Frequency Change ☐ Order Renewal

Is this the first dose? ☐ Yes ☐ No, date of last infusion: _____ Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

DIAGNOSIS AND ICD-10 CODE

☐ Rheumatoid Arthritis (RA) ICD-10 Code: _____

☐ Polyarticular Juvenile Idiopathic Arthritis (pJIA) ICD-10 Code: _____

☐ Psoriatic Arthritis (PsA) ICD-10 Code: _____

☐ Other: _____ ICD-10 Code: _____

REQUIRED DOCUMENTATION

☐ This signed order form by the provider

☐ Patient demographics AND insurance information

☐ TB Test Results

☐ H&P and Clinical/Progress notes supporting primary diagnosis

☐ Labs and Tests supporting primary diagnosis

☐ HBV screening results

List Tried & Failed Therapies, including duration of treatment:

1) k
2) k
3) k

MEDICATION ORDERS

Dosing

☐ Orencia _____mg IV over 30 minutes at Week 0, 2, 4 then every 4 weeks (Max dose= 1000mg)

☐ Orencia _____mg IV over 30 minutes every 4 weeks (Max dose= 1000mg)

☐ Orencia _____mg IV over 30 minutes x 1 dose, then _____mg SubQ once weekly

☐ Orencia _____mg SubQ once weekly

Refills*: ☐ X 6 months ☐ X 1 Year ☐ _____ doses

*(if not indicated, order will expire 1 year from date signed)

■ RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline

■ RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration

■ RN to flush and lock VAD/CVAD per company protocol

Other: _____

PREMEDICATION ORDERS

☐ Acetaminophen 650mg PO prior to infusion

☐ Diphenhydramine 25mg PO prior to infusion

☐ Other: _____

☐ Other: _____

EMERGENCY MEDICATIONS

■ Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO
Diphenhydramine 25mg-50mg slow IV push over 2-5 mins
Acetaminophen 325mg-650mg PO
Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated
Epinephrine 0.3mg IM/SQ, may repeat x1
Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO
Diphenhydramine 25mg slow IV push over 2-5 mins
Acetaminophen 325mg PO
Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated
Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1
Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (AIC/AIS only)

PRESCRIBER INFORMATION

Prescriber Name: _____ NPI Number: _____

Office Phone: _____ Office Fax: _____

Prescriber Signature: _____ Date: _____

© Access Infusion Care
REV: 6/2025

Gaylord, MI
Phone: (989) 619-6253
Fax: (855) 857-3834

Traverse City, MI
Phone: (231) 252-0893
Fax: (855) 857-3834

Marquette, MI
Phone: (906) 661-1088
Fax: (855) 857-3834