

PATIENT INFORMATION

Name: _____ DOB: _____ Height: _____ Weight: _____
 Address: _____ Phone: _____ Allergies: _____
 City, State, Zip: _____ Email: _____

REFERRAL STATUS

New Referral Dose or Frequency Change Order Renewal

Is this the first dose? Yes No, date of last infusion: _____ Line type: PIV PICC Port Other

DIAGNOSIS AND ICD-10 CODE

Severe Eosinophilic Asthma ICD-10 Code: _____
 Other: _____ ICD-10 Code: _____

Does your patient have blood eosinophil counts \geq 150 cells/ μ L within past 12 months? Yes No

REQUIRED DOCUMENTATION

- This signed order form by the provider
- Patient demographics AND insurance information
- Pulmonary Function Tests
- H&P and Clinical/Progress notes supporting primary diagnosis
- Labs and Tests supporting primary diagnosis, including blood eosinophil counts

List Tried & Failed Therapies, including duration of treatment:

- 1)
- 2)
- 3)

MEDICATION ORDERS

Initial Dosing Fasenra _____mg SubQ every 4 weeks for three doses then every 8 weeks thereafter
 Maintenance Dosing Fasenra _____mg SubQ every 8 weeks

Refills*: X 6 months X 1 Year Other: _____
 *(if not indicated, order will expire 1 year from date signed)

- RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline
 - RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration
 - RN to flush and lock VAD/CVAD per company protocol
- Other: _____

PREMEDICATION ORDERS

Acetaminophen 650mg PO prior to infusion Other: _____
 Diphenhydramine 25mg PO prior to infusion Other: _____

EMERGENCY MEDICATIONS

Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO
 Diphenhydramine 25mg-50mg slow IV push over 2-5 mins
 Acetaminophen 325mg-650mg PO
 Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.3mg IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive
 Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (AIC/AIS only)

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO
 Diphenhydramine 25mg slow IV push over 2-5 mins
 Acetaminophen 325mg PO
 Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

PRESCRIBER INFORMATION

Prescriber Name: _____ NPI Number: _____
 Office Phone: _____ Office Fax: _____
 Prescriber Signature: _____ Date: _____